

Product Name: GP Mousemaster 25  
APVMA Approval No: 93312/138705



Label Name:	GP Mousemaster 25
-------------	-------------------

Signal Headings:	DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	ACTIVE CONSTITUENT: 25g/kg ZINC PHOSPHIDE
-------------------------	---

Mode of Action:	
-----------------	--

Statement of Claims:	For the control of heavy infestations of mice in agricultural situations.  THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORIZED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.
----------------------	--

Net Contents:	12.5kg - 500kg
---------------	----------------

Restrains:	RESTRAINTS: DO NOT apply bait to bare ground (including fallow where there is no vegetative cover). DO NOT apply bait in a trail. DO NOT apply bait if heavy rain is imminent. DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals. DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.
------------	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.
Withholding Periods:	WITHHOLDING PERIOD DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.
Trade Advice:	
General Instructions:	
Resistance Warning:	
Precautions:	
Protections:	<p><b>PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK</b>  Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.</p>
Storage and Disposal:	<p><b>STORAGE AND DISPOSAL</b>  The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Excess or unused baits must be buried below one metre. Do not burn empty containers or product.</p>
Safety Directions:	<p><b>SAFETY DIRECTIONS</b>  Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits wear elbow length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber wash with detergent and warm water</p>
First Aid Instructions:	FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Do not give direct mouth to mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy viva or one way mask. Resuscitate in a well ventilated area.

First Aid Warnings:

## DIRECTIONS FOR USE

Situation	Pest	Rate	Critical Comments
<p>Grain crops, legume crops, canola, safflower, nut crops and pasture.</p> <p>(Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).</p>	Mice	1kg/ha to achieve an even coverage of 1 pellet/m <sup>2</sup>	<p>Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 1 pellet/m<sup>2</sup>. It is imperative that all attempts are made to apply bait as evenly as possible to maximize product efficacy and minimise risks to non-target animals. GP MOUSEMASTER 25 will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances.</p> <p>It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying the bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.</p>

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.